



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

better kind is \$1 and of the inferior \$.75 per lb. Now, he takes 9 parts of the better and mixes it with 2 parts of the inferior, then 9 parts of the mixture with 2 parts of the inferior, etc. What is the price of the  $n$ th mixture per lb.?

22. Proposed by F. P. MATZ, M. A., M. Sc., Ph. D., Editor of the Department of Mathematics in the "New England Journal of Education" and Professor of Mathematics and Astronomy in New Windsor College, New Windsor, Maryland.

For the sum  $D = \$30$ , Messrs. Zerr and Ellwood contracted to plough the sod for a circular track, width  $m = 60$  feet and inner radius  $r = 940$  feet. How is the money to be divided, if they commence ploughing at the inner circumference of the track, make uniform furrows of width  $n = 1\frac{1}{2}$  feet, and Mr. Ellwood continually follows Mr. Zerr during the ploughing?

Solutions to these problems should be received on or before July 1st.

---

## GEOMETRY.

---

Conducted by B. F. FINKEL, Kidder, Mo. All contributions to this department should be sent to him.

---

### SOLUTIONS TO PROBLEMS.

---

6. Proposed by EARL D. WEST, West Middleburg, Logan County, Ohio.

Having the sides 6, 4, 5, and 3 respectively of a trapezium, inscribed in a circle, to find the diameter of the circle.

- II. Solution by Professor G. B. M. ZERR, A. M., Principal of High School, Staunton, Virginia; and J. A. CALDERHEAD, Superintendent of Schools, Lima, Ohio; and CHARLES E. MYERS, Canton, Ohio.

Let  $ABCD$  be the quadrilateral inscribed in a circle,  $AC$  the diagonal,  $AB = a = 5$ ,  $BC = b = 3$ ,  $DC = c = 6$ , and  $DA = d = 4$ .

From the triangles  $ABC$  and  $CDA$

$$AC^2 = a^2 + b^2 - 2ab \cos B$$

$$AC^2 = c^2 + d^2 - 2cd \cos D = c^2 + d^2 + 2cd \cos B$$

$$\therefore \cos B = \frac{a^2 + b^2 - c^2 - d^2}{2(ab + cd)}.$$

$$\sin^2 B = 1 - \frac{(a^2 + b^2 - c^2 - d^2)^2}{4(ab + cd)^2}$$

$$= \frac{(c + b + d - a)(a + b + d - c)(a + c + d - b)(a + b + c - d)}{4(ab + cd)^2}$$

$$= \frac{16(s - a)(s - b)(s - c)(s - d)}{4(ab + cd)^2} \quad \text{where } s = \frac{1}{2}(a + b + c + d)$$

$$AC^2 = c^2 + d^2 + \frac{2cd(a^2 + b^2 - c^2 - d^2)}{2(ab + cd)} = \frac{(ac + bd)(ad + bc)}{(ab + cd)}.$$

$$R = \text{radius} = \frac{AC}{2 \sin b} = \frac{1}{4} \sqrt{\frac{(ab+cd)(ac+bd)(ad+bc)}{(s-a)(s-b)(s-c)(s-d)}}$$

$$2R = D = \frac{1}{2} \sqrt{\frac{(39)(42)(38)}{(4)(6)(3)(5)}} = \frac{1}{2} \sqrt{172.9} = 6.57457225.$$

Also solved by *P. S. BERG, H. C. WHITAKER, J. R. BALDWIN, P. H. PHILBRICK, J. W. SCHEFFER.*

[Note.—The formula for the area of an inscriptable quadrilateral is

$$A = \sqrt{s(s-a)(s-b)(s-c)(s-d)}, \text{ where } s = \frac{1}{2}(a+b+c+d). \text{—ED.}]$$

7. Proposed by **WILLIAM HOOVER, A. M., Ph. D.,** Professor of Mathematics and Astronomy in the Ohio University, Athens, Ohio.

Through each point of the straight line  $x=my+h$  is drawn a chord of the parabola  $y^2=4ax$ , which is bisected in the point. Prove that this chord touches the parabola  $(y+2am)^2=8a(x-h)$ .

II. Solution by **L. E. PRATT, Tecumseh, Nebraska.**

Let  $y=Bx+C \dots (1)$  be a straight line cutting the given parabola. The co-ordinates of the middle point of the chord intercepted by the curve are

$$\left( \frac{2a-BC}{B^2}, \frac{2a}{B} \right).$$

Substituting these for  $x$  and  $y$  in the equations of the given straight line, we have

$$BC=2a-2amB-hB^2 \dots (2).$$

If  $(x_1, y_1)$  be any point of (1) we have  $y_1=Bx_1+C \dots (3)$ .

Eliminating  $C$  from (2) and (3) we obtain a quadratic in  $B$  which may be written

$$B = \frac{y_1 + 2am \pm \sqrt{(y_1 + 2am)^2 - 8a(x_1 - h)}}{2(x_1 - h)} \dots (4).$$

This result shows that two chords bisected by the straight line  $x=my+h$  may be drawn through the point  $x_1, y_1$ ; that when the roots of  $B$ , the angular coefficient, are equal the two chords coincide in one; that this takes place when the radical in (4) is equal to zero. But when the radical equals zero the point  $x_1, y_1$ , is a point of the parabola

$$(y+2am)^2=8a(x-h)$$

and the straight line (1) is tangent to it.

This problem was solved in a very excellent manner by *Professors HUME, SCHEFFER, and ZERR.*

9. Proposed by **J. C. GREGG, Superintendent of Schools, Brazil, Indiana.**

Two circles intersect in  $A$  and  $B$ . Through  $A$  two lines  $CAE$  and  $DAF$  are drawn, each passing through a centre and terminated by the circumference. Show that  $CA \times AE = DA \times AF$ . [*Euclid.*]

Solution by **Miss GRACE H. GRIDLEY, Student in Kidder Institute, Kidder, Missouri.**

Let the straight lines  $CAE$  and  $DAF$  pass through the point of intersection  $A$  of the two circles, and through the centers  $O$  and  $O'$ , respectively.